

Table of Contents

7	Human Health	1
7.1	Environmental Setting	1
7.1.1	Population Characteristics of the Program Area.....	1
7.1.2	Hazards, Toxicity, and Exposure in the Environmental Setting.....	2
7.1.3	Pesticides and the Environment	3
7.1.4	Regulatory Environment	4
7.2	Environmental Impacts and Mitigation Measures	7
7.2.1	Evaluation Concerns and Criteria	7
7.2.2	Evaluation Methods and Assumptions.....	9
7.2.3	Surveillance Alternative	12
7.2.4	Physical Control Alternative	12
7.2.5	Vegetation Management Alternative.....	13
7.2.6	Biological Control Alternative	16
7.2.7	Chemical Control Alternative	16
7.2.8	Other Nonchemical Control/Trapping Alternative	24
7.2.9	Cumulative Impacts.....	25
7.2.10	Environmental Impacts Summary.....	25
7.2.11	Mitigation and Monitoring.....	29

Tables

Table 7-1a	Population and Growth in Contra Costa County (1990–2010)	1
Table 7-1b	Population and Growth in the Four Counties Adjacent to the Districts (1990–2010)	2
Table 7-2	Pesticide Active Ingredients.....	4
Table 7-3	Herbicide Active Ingredients and Adjuvants	4
Table 7-4	Herbicides Potentially Used for Mosquito Abatement.....	13
Table 7-5	Adjuvants Used for Weed Abatement.....	15
Table 7-6	Active Ingredients Identified for Further Evaluation in Appendix B.....	17
Table 7-7	Chemicals Employed for Larval Mosquito Abatement.....	17
Table 7-8	Chemicals Employed for Adult Insect Abatement.....	20
Table 7-9	Chemicals Employed for Rodent Abatement.....	23
Table 7-10	Summary of Alternative Human Health Impacts.....	27

This page left intentionally blank

7 Human Health

Chapter 7 evaluates potential impacts of the Program on human health. Results of the evaluation are provided at the programmatic level. Section 7.1, Environmental Setting, presents an overview of the District's human population and growth estimates and the federal and state regulations that are applicable to the Program. Section 7.2, Environmental Impacts and Mitigation Measures, presents the following:

- > Environmental concerns and evaluation criteria: A discussion of whether the Program alternatives would cause any potentially significant impacts to human health and addressing concerns from the public scoping
- > Discussion of methods and assumptions
- > Discussion of potential impacts of the Program alternatives, and mitigation measures, if necessary, for those impacts
- > Cumulative impacts summary
- > A summary of estimated impacts to human health
- > Potential ecological impacts are addressed in Chapter 6.

7.1 Environmental Setting

The Program Area is defined as Contra Costa Mosquito and Vector Control District (CCMVCD, the District) and surrounding counties including Alameda, Sacramento, San Joaquin and Solano counties that are impacted by unwanted vectors that must be managed to minimize adverse effects, disease, and environmental effects. The following section provides background information on the environmental fate and toxicity of pesticides and an overview of the regulatory setting with respect to chemical and biological pesticides.

7.1.1 Population Characteristics of the Program Area

The size of the population in the District's Service Area and the larger Program Area are shown in the following two tables. In 2010, the population of California was estimated at 37.3 million (US Census Bureau, (<http://www.bayareacensus.ca.gov/counties/counties.htm>)). The population of the nine counties in the nine Districts' Service Areas is approximately 6.76 million, which represents 18 percent of the statewide total (see Table 7-1a).

Table 7-1a Population and Growth in Contra Costa County (1990–2010)

County / Area	Population			Population Growth (Compound Annual Average)	
	1990	2000	2010	1990–2000	2000–2010
Contra Costa	803,732	948,816	1,049,025	1.67%	1.01%
Statewide Area	29,760,021	33,871,648	37,253,956	1.30%	0.96%

Table 7-1b provides the population counts and projected growth in the four counties adjacent to the Districts' Service Areas that are included in the Districts' Program Area.¹

Table 7-1b Population and Growth in the Four Counties Adjacent to the Districts (1990–2010)

County / Area	Population			Population Growth (Compound Annual Average)	
	1990	2000	2010	1990–2000	2000–2010
Alameda	1,279,182	1,443,741	1,510,271	1.22%	0.45%
Sacramento	1,066,789	1,229,940	1,418,788	1.43%	1.44%
San Joaquin	480,628	567,885	685,306	1.68%	1.90%
Solano	340,421	394,542	413,344	1.49%	0.47%
Adjacent County Total	3,167,020	3,636,108	4,027,709	1.39%	1.03%
Statewide Total	29,760,021	33,871,648	37,253,956	1.30%	0.96%

The California Department of Finance projects steady population growth in the future, with total state population reaching over 44 million by 2030. These projections represent a compound annual average population growth of 0.86 percent.

7.1.2 Hazards, Toxicity, and Exposure in the Environmental Setting

A “hazardous material” is defined in California Health and Safety Code Section 25501 (p): as “any material that, because of its quantity, concentration, or physical or chemical characteristics, poses a significant present or potential hazard to human health and safety or to the environment if released into the workplace or the environment. “Hazardous materials” include, but are not limited to, “hazardous substances, hazardous waste, and any material that a handler or the administering agency has a reasonable basis for believing that it would be injurious to the health and safety of persons or harmful to the environment if released into the workplace or the environment.” Any liquid, solid, gas, sludge, synthetic product, or commodity that exhibits characteristics of toxicity, ignitability, corrosiveness, or reactivity has the potential to be considered a “hazardous material.”

7.1.2.1 Toxicity and Exposure

Toxicology is the study of a compound’s potential to elicit an adverse effect in an organism. The toxicity of a compound is dependent upon exposure, including the specific amount of the compound that reaches an organism’s tissues (i.e., the dose), the duration of time over which a dose is received, the potency of the chemical for eliciting a toxic effect (i.e., the response), and the sensitivity of the organism receiving the dose of the chemical. Toxicity effects are measured in controlled laboratory tests on a dose/response scale, whereby the probability of a toxic response increases as dose increases. Exposure to a compound is necessary for potential toxic effects to occur. However, exposure does not, in itself, imply that toxicity will occur. Thus, toxic hazards can be mitigated by limiting potential exposure to ensure that doses are less than the amount that may result in adverse health effects.

The toxicity data included in the numerous tables and charts in this document are generally derived from rigidly controlled laboratory animal studies designed to determine the potential adverse effects of the chemical under several possible routes of exposure. In these studies, the species of interest is exposed to 100 percent chemical at several doses to determine useful information such as the lowest concentration resulting in a predetermined adverse effect (LOAEL) on numerous selected physiological and behavioral systems. The second component of these tests is to determine the highest concentration of chemical that results in no measurable adverse effect (NOAEL).

However, these, and other, coordinated and focused laboratory tests are designed to document the effects of the chemical when a continuous, controlled, exposure exists and do not realistically reflect the likely exposures or toxicity in the District field application scenarios. As such, the toxicity information is

intended as an overview of potential issues and guidance for understanding the completely “safe” maximum exposure levels of applications that would not adversely impact humans or nontarget plant and animal species.

Although the regulatory community uses this basic information to provide a relative comparison of the potential for a chemical to result in unwanted adverse effects and this information is reflected in the approved usage labels and MSDSs, in actual practice, the amounts applied in the District’s Program Area are often substantially less than the amounts used in the laboratory toxicity studies. Because of these large safety factors used to develop recommended product label application rates, the amount of chemical resulting in demonstrated toxicity in the laboratory is much higher than the low exposure levels associated with an actual application. The application concentrations consistent with the labels or MSDSs are designed to be protective of the health of humans and other nontarget species (i.e., low enough to not kill them, weaken them, or cause them to fail to reproduce). However, adverse effects may still occur to some nontarget organisms.

Although laboratory toxicity testing focuses on tiered concentrations of chemical exposure, the results of these tests produce a series of toxicity estimates of concentrations less than those that produce mortality. Extrapolation of these data is used to generate estimates of chronic toxicity or possible effects of lower doses that may result in sublethal effects such as reproduction or metabolic changes. In reality, these low-dose exposures need to be sustained over longer periods than are relevant to typical application scenarios for vector control including multiple applications in an area such as a wetland.

7.1.2.2 Chemistry, Fate, and Transport

Various biological, chemical, and physical parameters affect the behavior of a compound in the environment and its potential toxicity. The chemistry, fate, and transport of a compound must be analyzed to fully estimate potential exposure. The fate and transport of a compound is determined by the physical and chemical properties of the compound itself and the environment in which it is released. Thus, the following characteristics of a compound must be evaluated: its half-life in various environmental media (e.g., sediment, water, air); photolytic half-life; lipid and water solubility; adsorption to sediments and plants; and volatilization. Environmental factors that affect fate and transport processes include temperature, rainfall, wind, sunlight, water turbidity, and water and soil pH. Information pertaining to these parameters allows evaluation of how compounds may be transported between environmental media (e.g., from sediments to biota), how a compound may be degraded into various breakdown products, and how long a compound or its breakdown products may persist in different environmental media. Appendix B, Ecological and Human Health Risk Assessment, provides a discussion of the environmental fate of the pesticide active ingredients and other chemicals associated with specific pesticide formulations used in the Program alternatives.

7.1.3 Pesticides and the Environment

The pesticide and herbicide active ingredients included in the Proposed Program are listed in Table 7-2 and Table 7-3, respectively. Appendix B provides results of review and evaluations of the active ingredients and adjuvants the District currently uses or proposes to use

Table 7-2 Pesticide Active Ingredients

Active Ingredient	Vector
Bacillus sphaericus (Bs)	Mosquito (larvae)
Bacillus thuringiensis israelensis (Bti)	Mosquito (larvae)
Spinosad	Mosquito (larvae)
Biodegradable alcohol ethoxylated surfactant	Mosquito (larvae and pupae)
Methoprene	Mosquito (larvae)
Mineral Oil	Mosquito (larvae and pupae)
Petroleum distillate	Mosquito (larvae and pupae)
Temephos	Mosquito (larvae)
Pyrethrins	Mosquito (adult), Yellow jacket wasp
Piperonyl butoxide (PBO)	Mosquito (adult), Yellow jacket wasp
Naled	Mosquito (adult)
Resmethrin	Mosquito (adult)
Permethrin	Mosquito (adult), Tick
Etofenprox	Mosquito (adult)
Diphacinone	Rat
Bromadiolone	Rat
Difethialone	Rat
Cholecalciferol	Rat

Table 7-3 Herbicide Active Ingredients and Adjuvants

Active Ingredient/Adjuvant	Vector
Glyphosate	Weeds
2,4-DP (dimethylamine salt)	Weeds

7.1.4 Regulatory Environment

Formulations proposed for each Program alternative for vector control are and would be used according to federal and state regulatory requirements for the registration, transportation, and use of pesticides. The regulatory framework pertaining to the use of pesticides is discussed below.

7.1.4.1 Federal

The USEPA regulates pesticides under two major statutes: FIFRA and Federal Food, Drug, and Cosmetic Act. Under these acts, the USEPA mandates extensive scientific research to assess risks to humans, domestic animals, wildlife, plants, groundwater, and beneficial insects before granting registration for a pesticide. These studies allow the USEPA to assess the potential for human and ecological health effects. When new data raise concern about a registered pesticide's safety, the USEPA may take action to suspend or cancel its registration. The USEPA may also perform an extensive special review of a pesticide's risks and benefits and/or work with manufacturers and users to implement changes in a pesticide's approved use (e.g., reducing application rates).

7.1.4.1.1 Federal Insecticide, Fungicide, and Rodenticide Act

FIFRA defines a pesticide as "any substance intended for preventing, destroying, repelling, or mitigating any pest." The act requires USEPA registration of pesticides prior to their distribution for use in the US, sets registration criteria (testing guidelines), and mandates that pesticides perform their intended functions without causing unreasonable adverse effects on people and the environment when used according to USEPA-approved label directions. FIFRA defines an "unreasonable adverse effect on the environment" as "(1) any unreasonable risk to man or the environment, taking into account the economic, social, and environmental costs and benefits of the use of the pesticide, or (2) a human dietary risk from residues that result from a use of a pesticide in or on any food inconsistent with the standard under Section 408 of the Federal Food, Drug, and Cosmetic Act (21 USC 346a)."

FIFRA regulates only the active ingredients of pesticides, not inert ingredients, which manufacturers are not required to reveal. However, toxicity studies conducted under FIFRA are required to evaluate the active ingredient and the entire product formulation, through which any potential additive or synergistic effects of inert ingredients are established.

7.1.4.1.2 Federal Food, Drug, and Cosmetic Act

This act authorizes the USEPA to set tolerances (i.e., maximum allowable amounts) for pesticide residues in/on food. Thus, this act does not expressly regulate pesticide use, but exceedance of tolerances may result in prosecution or changes in the approved use of a pesticide regulated under FIFRA.

7.1.4.1.3 Clean Water Act and National Pollutant Discharge Elimination System

The CWA establishes the principal federal statutes for water quality protection "to restore and maintain the chemical, physical, and biological integrity of the nation's water, to achieve a level of water quality which provides for recreation in and on the water, and for the propagation of fish and wildlife."

- > Section 303(d) requires each state to provide a list of impaired waters that do not meet or are expected not to meet state water quality standards as defined by that section. The CWA regulates potentially toxic discharges through the NPDES and ambient water quality through numeric and narrative water quality standards. The release of aquatic pesticides into waters of any state may require an NPDES permit, depending on the pesticide considered, and the conditions proposed for application.
- > Section 402, the NPDES, requires permits for pollution discharges (except dredge or fill material) into US waters, such that the permitted discharge does not cause a violation of federal and state water quality standards. Biological and residual pesticides discharged into surface waters constitute pollutants and require coverage under an NPDES permit. In California, NPDES permits are issued by the SWRCB or the RWQCBs.

7.1.4.1.4 Safe Drinking Water Act of 1974

Under the Safe Drinking Water Act of 1974, the USEPA establishes Maximum Contaminant Levels (MCLs), which are specific concentrations that cannot be exceeded for a given contaminant in surface

water or groundwater. USEPA has the ability to enforce these nationwide standards or delegate administration and enforcement duties to state agencies. The CDPH administers the federal Safe Drinking Water Act in California.

7.1.4.1.5 California Toxics Rule

In 2000, the USEPA developed water quality criteria for priority toxic pollutants to protect human health and the environment when a gap in California's water quality standards was created when the state's water quality control plans containing water quality criteria for priority toxic pollutants were overturned in 1994 (thus causing California to be out of compliance with the CWA). These established criteria are to be applied to inland surface waters, enclosed bays, and estuaries in California. The rule includes aquatic life criteria for 23 priority toxic pollutants, human health criteria for 57 priority toxics, and a compliance schedule.

7.1.4.2 State of California

California's programs for the registration of pesticides and commercial chemicals parallel federal programs, but many of California's requirements are stricter than federal requirements. The Cal/EPA regulates registration of pesticides and commercial chemicals in California. Within Cal/EPA, the CDPR oversees pesticide evaluation and registration through use enforcement, environmental monitoring, residue testing, and reevaluation. The CDPR works with County Agricultural Commissioners, who evaluate, develop conditions of use, approve, or deny permits for restricted-use pesticides; certify private applicators; conduct compliance inspections; and take formal compliance or enforcement actions. The Secretary of Resources has certified California's pesticide regulatory program as meeting CEQA requirements (CDPR 2006).

California also requires commercial growers and pesticide applicators to report commercial pesticide applications to local county agricultural commissioners. The CDPR compiles this information in annual pesticide use reports. The CDPR's Environmental Hazards Assessment Program collects and analyzes environmental pesticide residue data, characterizes drift and other offsite pesticide movement, and evaluates the effect of application methods on movement of pesticides in air. If a pesticide is determined to be a toxic air contaminant, appropriate control measures are developed with the California Air Resources Board to reduce emissions to levels that adequately protect public health. Control measures may include product label amendments, applicator training, restrictions on use patterns or locations, and product cancellations.

7.1.4.2.1 Porter-Cologne Act and State NPDES Permitting

Under the Porter-Cologne Act (California Water Code Section 13000) the SWRCB, and the state's nine RWQCBs that it oversees, are responsible for administering federal and state water quality regulation and permitting duties.

The SWRCB oversees pesticide NPDES permitting in California. Users of specific larvicide and adulticide registered products are required to obtain coverage under the Statewide NPDES Permit for Biological and Residual Pesticide Discharges to Waters of the US from Vector Control Applications (SWRCB Water Quality Order No. 2012-0003-DWQ; NPDES No. CAG 990004; Vector Control Permit). Users of certain aquatic herbicides are required to obtain coverage under the Statewide General NPDES Permit for the Discharge of Aquatic Pesticides for Aquatic Weed Control in Waters of the US (SWRCB Water Quality Order No. 2004-0009-DWQ; NPDES No. CAG 990005; Aquatic Weed Control Permit). Pesticides and herbicides that require state NPDES permitting include Bti, Bs, temephos, spinosad, petroleum distillates, naled, pyrethrin, permethrin, resmethrin, prallethrin, PBO, etofenprox, 2,4-D, glyphosate, imazapyr, and triclopyr. Both permits are discussed in detail in Chapter 9, Section 9.1.2.2.9.

7.1.4.2.2 Safe Drinking Water Act 1976

The CDPH administers the federal Safe Drinking Water Act in California. In addition to enforcing the primary MCLs (discussed above in Section 7.1.4.1), CDPH uses as guidelines Secondary MCLs that regulate constituents that affect water quality aesthetics (such as taste, odor, or color).

Additionally, under the California Safe Drinking Water Act, Cal/EPA's Office of Environmental Health Hazard Assessment develops Public Health Goals (PHGs) for contaminants in California's publicly supplied drinking water. PHGs are concentrations of drinking water contaminants that pose no significant health risk if consumed for a lifetime, based on current risk assessment principles, practices, and methods. Public water systems use PHGs to provide information about drinking water contaminants in their annual Consumer Confidence Reports.

7.1.4.2.3 The Safe Drinking Water and Toxic Enforcement Act (Proposition 65)

This act, passed as a ballot initiative in 1986, requires the state to annually publish a list of chemicals known to the state to cause cancer or reproductive toxicity so that the public and workers are informed about exposures to potentially harmful compounds. Cal/EPA's Office of Environmental Health Hazard Assessment administers the act and evaluates additions of new substances to the list. Proposition 65 requires companies to notify the public about chemicals in the products they sell or release into the environment, such as through warning labels on products or signs in affected areas, and prohibits them from knowingly releasing significant amounts of listed chemicals into drinking water sources.

7.1.4.2.4 California Pesticide Regulatory Program

CDPR regulates the sale and use of pesticides in California. CDPR is responsible for reviewing the toxic effects of pesticide formulations and determining whether a pesticide is suitable for use in California through a registration process. Although CDPR cannot require manufacturers to make changes in labels, it can refuse to register products in California unless manufacturers address unmitigated hazards by amending the pesticide label. Consequently, many pesticide labels that are already approved by the USEPA also contain California-specific requirements. Pesticide labels defining the registered applications and uses of a chemical are mandated by USEPA as a condition of registration. The label includes instructions telling users how to make sure the product is applied only to intended target pests, and includes precautions the applicator should take to protect human health and the environment. For example, product labels may contain such measures as restrictions in certain land uses and weather (i.e., wind speed) parameters.

7.2 Environmental Impacts and Mitigation Measures

This section evaluates the potential impacts from the Program alternatives, focusing on the human health impacts specific to the use of selected chemical and biological pesticides.

7.2.1 Evaluation Concerns and Criteria

The public has indicated concerns about some of the following issues. While not required, the responses to the concerns help to direct the reader to the appropriate section or Appendix B, Ecological and Human Health Risk Assessment, or they provide explanatory information in concise form.

- > The PEIR should address Program impacts on people and pets through ingestion and absorption pathways and proposed mitigation. Address impacts on chemically sensitive people and sensitive populations such as children, the elderly, pregnant women. Exposure to pesticides can result in compromised immune system, which would allow for development of allergies or autoimmune disorders.

- Potential Chemical Control Alternative impacts are discussed in Section 7.2.7, and toxicity of individual active ingredients is evaluated in greater detail in Appendix B.
- > The PEIR must list any and all biological or chemical agents proposed for use.
 - The biological and chemical pesticide formulations included in the Program are listed in Table 7-2, Pesticide Active Ingredients and Table 7-3, Herbicide Active Ingredients and Adjuvants.
- > CDPH should be consulted to ensure all potential risks are identified, characterized, and evaluated.
 - The PEIR document and information will be made publicly available and will be reviewed by the appropriate regulatory bodies.
- > Concern expressed over public safety and health with regards to existing vegetable gardens and fruit trees within the project area. Local swimming holes could be a potential habitat for breeding mosquitoes, and chemical treatment could impact humans.
 - BMPs to reduce exposure to nontarget species and areas are discussed in Chapter 2, Section 7.2.7 herein, summarized in several other relevant chapters, and evaluated in greater detail in Appendix B.
- > Concerned with use of Zenivex; it mimics chrysanthemums but is a harmful neurotoxin.
 - Etofenprox, the active ingredient in Zenivex, is discussed in Section 7.2.7.2.2 and evaluated in greater detail in Appendix B. It does not require concomitant use of a synergist, such as PBO. Therefore, it likely exhibits less toxicity than others that require co-application with other chemicals. Based on toxicity, environmental fate, and usage patterns, etofenprox is not likely to result in unwanted adverse impacts to humans when BMPs are used.
- > Concerned that adulticides present danger to humans, as many are known carcinogens and endocrine disruptors.
 - The District's BMPs provide that adulticides are generally applied as aerosols using ULV techniques to minimize exposure to nontarget species. Aerial and ground application techniques are used to distribute the insecticides. The potential toxicity of the various adulticides included in the Program are discussed in Section 7.2.7 and evaluated in greater detail in Appendix B.
- > Concerned that pyrethrins disrupt the normal functioning of sex hormones, while PBO affects the functioning of hormone-related organs.
 - The District generally uses pyrethrins in ULV applications, which are designed to prevent environmental persistence and potential impacts to nontarget species.
 - As a synergist for pyrethrins and pyrethroids, PBO is also generally applied by ULV, and it degrades rapidly in soil and water. Its potential toxicity is discussed in Appendix B.
- > In addition to short-term effects, what are the long-term effects of repeated exposure to these chemicals?
 - The chronic effects of the various pesticides are discussed in detail in Appendix B.

The CEQA Guidelines Appendix G, Environmental Checklist Form, does not contain criteria for determining significance of impacts to human health from the use of pesticides and herbicides. The criteria for hazards and hazardous materials (Checklist Section VIII) are primarily addressed in Chapter 8. However, the first criterion is partly applicable and asks would the project:

- a. Create a significant hazard to the public or the environment through the routine transport, use, or disposal of hazardous materials?

The applicability is for the use of these chemicals. In short, the determination of significance is based on the potential to adversely affect human health based on existing data and application methods including label requirements and additional BMPs the District employs (see Section 2.9). The specific concern is whether the activities used to control pest species could result in direct or indirect impacts to human populations in a treatment area in the short term (i.e., acute toxicity) or over the long term (i.e., chronic toxicity).

7.2.2 Evaluation Methods and Assumptions

Pesticides the District uses were investigated to provide a preliminary assessment of the potential impacts to humans (discussed in detail in Appendix B). A comprehensive literature review was conducted and the District supplied information to assess potential exposure and toxicity using the following:

- > Pesticides the District uses
- > Pesticide label recommendations
- > Types of application sites (e.g., habitat types)
- > Application procedures
- > Number of treatments per application site
- > Total amount used per treatment for each application site, based on yearly totals.
- > Physicochemical properties of the pesticides/active ingredients
- > Pesticide target vector efficacy
- > Reported adverse effects (e.g., reproductive, developmental, carcinogenic)

The pesticide application scenarios that result in reasonable efficacy with minimal unwanted estimated risk are preferred and are the basis of IPM/IVM approaches and BMPs the District practices. Each of the pesticides identified as warranting further evaluation in Appendix B are known to exhibit at least one parameter that appears to drive potential or perceived risk. Toxicity levels (e.g., slight, low, moderate, high, etc.) are used prevalently in the published literature but are not standardized or representative of specific criteria. They qualitatively describe toxicity in relative terms in the evaluations of herbicides and pesticides in this PEIR and in Appendix B. Toxicity levels are helpful in making significance determinations.

The pesticide application scenarios that result in reasonable efficacy with minimal unwanted estimated risk are preferred and are the basis of IPM/IVM approaches and BMPs the District employs. All BMPs are described in Chapter 2 (Table 2-9), and the most relevant BMPs for avoidance or minimization of impacts to human health are repeated below.

For all six Program alternatives, the District uses the following BMPs:

- > Operation of noise-generating equipment (e.g., chainsaws, wood chippers, brush-cutters, pickup trucks) will abide by the time-of-day restrictions established by the applicable local jurisdiction (i.e., City and/or County) if such noise activities would be audible to receptors (e.g., residential land uses, schools, hospitals, places of worship) located in the applicable local jurisdiction. Shut down all motorized equipment when not in use. (Table 2-9, BMP A11)
- > For operations that generate noise expected to be of concern to the public, the following measures will be implemented: (Table 2-9, BMP A12)

- Measure 1: Provide Advance Notices: A variety of measures are implemented depending on the magnitude/nature of the activities the District undertakes and may include, but are not limited to, press releases, hand-delivered flyers, and posted signs. Public agencies and elected officials also may be notified of the nature and duration of the activities, including the Board of Supervisors or City Council, environmental health and agricultural agencies, emergency service providers, and airports.
 - Measure 2: Provide Mechanism to Address Complaints: District staff is available during regular business hours to respond to service calls and address concerns about nighttime operations.
- > The District will perform public education and outreach activities. (Table 2-9, BMP A13)
 - > To minimize air and GHG emissions, engine idling times will be minimized either by shutting equipment and vehicles off when not in use or reducing the maximum idling time to 5 minutes. Correct tire inflation will be maintained in accordance with manufacturer's specifications on wheeled equipment and vehicles to prevent excessive rolling resistance. All equipment and vehicles will be maintained and properly tuned in accordance with manufacturer's specifications. All equipment will be checked by a certified visible emissions evaluator if visible emissions are apparent to onsite staff. (Table 2-9, BMP A14)
 - > A hazardous spill plan will be developed, maintained, made available, and staff trained on implementation and notification for petroleum- based or other chemical-based materials prior to commencement of vector treatment activities. (Table 2-9, BMP I5)
 - > Equip all vehicles used in wildland areas with a shovel and a fire extinguisher at all times. (Table 2-9, BMP J1)
 - > Train employees on the safe use of equipment and machinery, including vehicle operation. (Table 2-9, BMP J2)
 - > District will regularly review and update their existing health and safety plan to maintain compliance with all applicable standards. Employees will be required to review these materials annually. (Table 2-9, BMP J3)

For five Program alternatives, only excluding Biological Control, the following BMPs are protective of human health:

- > Vehicles driving on levees to travel through tidal marsh or to access sloughs or channels for surveillance or treatment activities will travel at speeds no greater than 10 miles per hour to minimize noise and dust disturbance. (Table 2-9, BMP A8)

For the Vegetation Management and Chemical Control Alternatives, the District uses the following BMPs:

- > District staff will conduct applications with strict adherence to product label directions that include approved application rates and methods, storage, transportation, mixing, and container disposal. (Table 2-9, BMP H1)
- > Materials will be applied at the lowest effective concentration for a specific set of vectors and environmental conditions. Application rates will never exceed the maximum label application rate.(Table 2-9, BMP H3)
- > To minimize application of pesticides, application of pesticides will be informed by surveillance and monitoring of vector populations. (Table 2-9, BMP J4)
- > District staff will follow label requirements for storage, loading, and mixing of pesticides and herbicides. Handle all mixing and transferring of herbicides within a contained area. (Table 2-9, BMP J5)
- > Postpone or cease application when predetermined weather parameters exceed product label specifications, when wind speeds exceed the velocity as stated on the product label, or when a high

chance of rain is predicted and rain is determining factor on the label of the material to be applied. (Table 2-9, BMP H6)

- > Applicators will remain aware of wind conditions prior to and during application events to minimize any possible unwanted drift to waterbodies, and other areas adjacent to the application areas. (Table 2-9, BMP H7)
- > Spray nozzles will be adjusted to produce larger droplet size rather than smaller droplet size. Use low nozzle pressures where possible (e.g., 30 to 70 pounds per square inch). Keep spray nozzles within a predetermined maximum distance of target weeds or pests (e.g., within 24 inches of vegetation during spraying). Adjusting droplet size would only apply to larvicides, herbicides, and non-ULV applications. Use ULV sprays that are calibrated to be effective and environmentally compatible at the proper droplet size (about 10-30 microns). (Table 2-9, BMP H8)
- > Clean containers at an approved site and dispose of at a legal dumpsite or recycle in accordance with manufacturer's instructions if available. (Table 2-9, BMP H9)
- > District staff will monitor sites post-treatment to determine if the target vector or weeds were effectively controlled with minimum effect to the environment and nontarget organisms. This information will be used to help design future treatment methods in the same season or future years to respond to changes in site conditions. (Table 2-9, BMP H11)
- > The District will provide notification to the public (24 to 48 hours in advance, if possible) and/or appropriate agency(ies) when applying pesticides or herbicides for large-scale treatments that will occur in close proximity to homes, heavily populated, high traffic, and sensitive areas. The District infrequently applies or participates in the application of herbicides in areas other than District facilities. (Table 2-9, BMP H13)
- > Exercise adequate caution to prevent spillage of pesticides during storage, transportation, mixing, or application of pesticides. Report all pesticide spills and cleanups (excepting cases where dry materials may be returned to the container or application equipment). (Table 2-9, BMP I1)
- > Maintain a pesticide spill cleanup kit and proper protective equipment at the District's Service Yard and in each vehicle used for pesticide application or transport. (Table 2-9, BMP I2)
- > Manage the spill site to prevent entry by unauthorized personnel. Contain and control the spill by stopping it from leaking or spreading to surrounding areas, cover dry spills with polyethylene or plastic tarpaulin, and absorb liquid spills with appropriate absorbent materials. (Table 2-9, BMP I3)
- > Properly secure the spilled material, label the bags with service container labels identifying the pesticide, and deliver them to a District Supervisor for disposal. (Table 2-9, BMP I4)
- > Field-based mixing and loading operations will occur in such a manner as to minimize the risk of accidental spill or release of pesticides. (Table 2-9, BMP I6)

This evaluation assumes that all pesticides are applied in accordance with label instructions and USEPA and Cal/EPA requirements. The USEPA requires mandatory statements to be included on pesticide product labels that include directions for use; precautions for avoiding certain dangerous actions; and where, when, and how the pesticide should be applied. This guidance is designed to ensure proper use of the pesticide and prevent unreasonable adverse effects to humans and the environment. All pesticide labels are required to include the name and percentage by weight of each active ingredient in the product/formulation. Toxicity categories for product hazards and appropriate first-aid measures must be properly and prominently displayed. Pesticide labels also outline proper use, storage, and disposal procedures, as well as precautions to protect applicators. The directions for use indicate target organism, appropriate application sites, application rates or dosages, contact times, and required application equipment for the pesticide. Warnings regarding appropriate wind speeds, droplet sizes, or habitats to avoid during application are also prominently displayed

This evaluation herein does not include assumptions about which alternative treatment strategy(ies) would be applied in any given area. Criteria used to trigger a particular alternative based on vector abundance and other variables are included in the District's operating procedures. This evaluation

assumes that important parameters, such as media half-life, are dependent on the specific conditions at the time of pesticide application, and values listed herein serve as reference values.

Concerning the application of multiple chemical treatments in the same area, such as larvicides followed by adulticides (i.e., not likely to occur under normal circumstances), or the application of multiple pesticides at the same time in a specific area (e.g., usually multiple active ingredients in the formulation such as VectoMax which combines Bti and Bs), the following information applies:

Most products sold as herbicides and pesticides are evaluated herein both for the active ingredient and for the adjuvants and surfactants used to make the product more useful. When multiple products are used in a vector control application, the impacts are weighed against the proximity and timing of each application. If products with an identical active ingredient are applied simultaneously, it is likely that the net effect could be the sum of the total active ingredient that is available for uptake by the vector. Although a synergy is possible in this scenario, it is typically not an approach used and is limited by the BMPs for that scenario. Because most pesticides and herbicides now have considerably less half-life (persistence), the overlap that would produce a residual exposure to a product would not occur unless the timing of applications is inappropriately close. Actual applications do not generally occur that close together unless there is a problem with treatment effectiveness. A material is applied followed by post treatment inspection to determine effectiveness. Only if the mosquitoes have not been sufficiently killed would the District go back into the area and reapply a pesticide.

7.2.3 Surveillance Alternative

Vector surveillance is critical to IVM strategies because it provides information that is used to determine when and where to institute other vector control measures. The District's mosquito surveillance activities are conducted in compliance with accepted federal and state guidelines (e.g., *California Mosquito-Borne Virus Surveillance and Response Plan* (CDPH et al. 2013) and *Best Management Practices for Mosquito Control in California* (CDPH and MVCAC 2012). These guidelines allow for flexibility in selection and specific application of control methods because local areas vary.

District practices would be a continuation of existing activities using applicable techniques, equipment, vehicles, and watercraft (except for possible purchase of an airboat for future use). Surveillance activities involve monitoring the distribution and abundance of adult and larval mosquitoes (field counting, sampling, and trapping), field inspection of mosquito habitat, testing for the presence of arboviruses in mosquitoes and their hosts, collection and testing of ticks for the presence of tickborne pathogens (e.g., lyme disease, ehrlichia, tularemia and spotted fever group rickettsia), small rodent trapping and testing, and/or response to public service requests regarding other vector animals or insects (e.g., yellow jacket wasps). Surveillance of potential areas of concern is a critical element for directing and responding to potential outbreaks of vectors and the potential for conveying vector-borne diseases.

Impact HH-1: No impact would occur to human health from the use of the Surveillance Alternative.

7.2.4 Physical Control Alternative

Physical Control for mosquitoes consists of the management of mosquito-producing habitat (including freshwater marshes and lakes, saltwater marshes, temporary standing water, and wastewater treatment facilities) especially through water control and maintenance or improvement of channels, tide gates, levees, and other water control facilities. Physical control is usually the most effective mosquito control technique because it provides a long-term solution by reducing or eliminating mosquito developmental sites and ultimately reduces the need for chemical applications. The physical control practices may be categorized into three groups: maintenance, new construction, and cultural practices. The District performs these physical control activities in accordance with all appropriate environmental regulations (wetland fill and dredge permits, endangered species review, water quality review, streambed alteration

permits, etc.), and in a manner that generally maintains or improves habitat values for desirable species. Physical control for other vectors such as rodents is based on the District’s site inspections to determine conditions promoting infestation, and property owners are provided educational materials on control measures that include removal of food sources and professionals to contact to remove the infestation. Physical control techniques have minimal impact on humans due to prior identification and avoidance of potential problem areas and wildlife habitats by publishing scheduled treatment times and locations.

Impact HH-2: Impacts to human health from use of the Physical Control Alternative would be **less than significant** and mitigation is not required.

7.2.5 Vegetation Management Alternative

The District uses hand tools (e.g., shovels, pruners, chainsaws, and weed-whackers) and heavy equipment where necessary for vegetation removal or thinning and sometimes applies herbicides to improve surveillance or reduce vector habitats. Vegetation removal or thinning primarily occurs in aquatic habitats to assist with the control of mosquitoes and in terrestrial habitats to help with the control of other vectors. To reduce the potential for mosquito breeding associated with water retention and infiltration structures, District staff may systematically clear or trim weeds and other obstructing vegetation in wetlands and retention basins (or request the structures’ owners to perform this task). These tasks are performed in conjunction with discussions with appropriate agencies (including CDFW) about special status species, and their habitats that must be acknowledged and protected prior to these activities. Vegetation management is also performed to assist other agencies and landowners with the management of invasive/nonnative weeds. These actions are typically performed under the direction of the concerned agency, which also maintains any required permits. These activities are conducted during predetermined times of recreational inactivity to provides an additional measure of safety to the public.

Impact HH-3: No impact would occur to human health from the nonherbicide Vegetation Management Alternative.

7.2.5.1 Herbicides

The herbicides the District may use in the future would be applied in strict conformance with label requirements. Herbicides the District may use in the future are discussed in Appendix B, and those that have been identified for additional evaluation are described below.

The District may use herbicides to control vegetation in and around mosquito habitats to improve access needed for surveillance and to reduce potential habitat for mosquitoes. The herbicides are listed in Table 7-4 and discussed in the indicated sections of Appendix B.

Table 7-4 Herbicides Potentially Used for Mosquito Abatement

Active Ingredient	Appendix B
Imazapyr	Section 4.6.1
Glyphosate	Section 4.6.2
Triclopyr	Section 4.6.3
2,4-Dichlorophenoxy acetic acid (2,4-D)	Section 4.6.4
Sulfometuron Methyl	Section 4.6.5

Herbicides are typically classified into the following major categories: pre-emergent herbicides (applied to the soil to prevent seedlings from germinating and emerging); post-emergent herbicides (applied after seedlings have emerged and control actively growing plants via contact damage or systemic impacts); contact herbicides (cause physical injury to the plant upon contact); and systemic herbicides (damage the internal functioning of the plant). Herbicides included in the Program have diverse chemical structures, act through distinct modes of action, and exhibit varying levels of potential toxicity to humans. Many of the herbicides are typically nonselective and broad-spectrum and function by inhibiting growth but do so in a multitude of ways. Most of the herbicides are moderately persistent in soil and water (for each herbicide's half-life in soil and water, please refer to Appendix B). The following have been shown to exhibit no/low toxicity to humans: imazapyr (USEPA 2006a), triclopyr (USEPA 1998a), and sulfometuron methyl (USEPA 2008). The actual use and human exposure in the field are far less than tested in the laboratory, and much higher volumes (exposure) would be needed to result in toxicity.

Many of the herbicides are typically nonselective and broad-spectrum and generally function by inhibiting growth but do so in a multitude of ways. For example, sulfometuron methyl retards or stops root and shoot development. Herbicides used against annual broadleaf weeds are generally of the post-emergent variety, such as triclopyr and sulfometuron methyl. In addition, imazapyr is a systematic, nonselective, pre- and post-emergent herbicide used for a broad range of terrestrial and aquatic weeds. Glyphosate represents a commonly used herbicide for the control and elimination of grass weeds and sedges. Most of the herbicides are moderately persistent in soil and water (for each herbicide's half-life in soil and water, refer to Appendix B).

Impact HH-4: Impacts to human health from the use of 2,4 D would be **less than significant and mitigation is not required.**

Glyphosate and adjuvants were identified for further evaluation based on use patterns and toxicity (Appendix B) and discussed in further detail below.

7.2.5.1.1 Glyphosate

Glyphosate is a nonselective, post-emergent, and systemic herbicide that is the active ingredient (as an acid or salt) in Alligare, Aquamaster, Buccaneer, and Roundup® products. It is designed to target the shikimic acid pathway, which is specific to plants and some microorganisms; therefore, glyphosate is thought to have very low toxicity to mammals (USEPA 1993). The District strictly adheres to its BMPs and product label requirements when using Glyphosate. Every effort is also made to minimize treatments that could affect milkweed, a plant important to Monarch butterfly populations.

The USEPA classifies glyphosate as Category III for oral and dermal toxicity (USEPA 1993), and the isopropylamine and ammonium salts exhibit low toxicity to mammals via the oral and dermal routes. No scientific evidence indicates that glyphosate is carcinogenic or mutagenic (USEPA 1993). It is poorly biotransformed in rats and is excreted via feces and urine; neither the parent compound nor its major breakdown product bioaccumulates in animal tissue (Williams et al. 2000). It is further mediated by the techniques used for application. Typically, spraying does not occur when wind is 5 mph or greater or within 15 feet of a crop area or sensitive habitat.

Despite the apparent lack of toxicity to mammals, concerns have been raised about glyphosate's long-term developmental and reproductive effects. Although still in review, glyphosate is included in the final list of chemicals for screening under USEPA's Endocrine Disruptor Screening Program (USEPA 2009a). The issue of endocrine-disrupting compounds is a topic of current scientific concern and inquiry but insufficient information is available to reach any conclusion about significance or potential adverse impacts. Recently, the USEPA renewed the temporary approval of a glyphosate and 2-4-D combination product (Enlist-Duo) for use with weed vectors, indicating it has not received significant adverse data to negate the decision (USEPA 2014a). This situation is true of all "potential" endocrine-disrupting compounds. Claims of endocrine disruption by dozens of chemicals are still an active area of research. Only very high exposures to those chemicals suggested as endocrine disruptors can be shown to suggest

a linkage. Because this exposure linkage is so high, endocrine disruption in a human would require, if real, exposure to substantially higher levels of chemical than used for vector control.

It is likely that USEPA will provide an updated review of its potential risks in 2015; however, current data indicate that glyphosate is nontoxic to humans.

Impact HH-5: Impacts to human health from the use of glyphosate would be **less than significant** and mitigation is not required.

7.2.5.2 Adjuvants

An adjuvant is any compound that is added to a pesticide/herbicide formulation or tank mix to facilitate the mixing, application, or effectiveness of that product. FIFRA does not require testing and registration of adjuvants. As such, little information on their fate, transport, and toxicity exists, other than that provided by the manufacturer or published by the scientific community (Bakke 2007; Tu et al. 2001). CDPR does require the registration of adjuvants that are considered to increase the action of the pesticides with which they are used (Bakke 2007). The adjuvants the District employs and where they are discussed in detail in Appendix B and are listed in Table 7-5.

Table 7-5 Adjuvants Used for Weed Abatement

Active Ingredient	Appendix B
APEs	Section 4.7.1
Polydimethylsiloxane Fluids	Section 4.7.2
Modified Vegetable Oil and Methylated Seed Oil	Section 4.7.3

APEs are used as detergents, dispersants, emulsifiers, solubilizers, and foaming and wetting agents. Primary degradation of APEs in the environment generates more persistent shorter chain compounds, some of which may mimic natural hormones and disrupt endocrine function in humans (Ying et al. 2002). Nonylphenol and nonylphenol ethoxylate, which are produced in large volumes and widely used, exhibit low acute oral and dermal toxicity but are highly irritating and corrosive to the skin and eyes (USEPA 2010). The acute toxicity of APEs to mammals is low; however, concern exists regarding the estrogen-mimicking behaviors of these compounds, particularly nonylphenol and nonylphenol ethoxylate (USEPA 2010). The USEPA (2010) has recently recommended that this suite of chemicals be evaluated further due to their widespread use, persistence, and possible estrogen-mimicking behavior.

Vegetable-derived oils (from soybeans, cottonseeds, etc.) decrease surface tension, but they are not as effective as other surfactants at increasing spreading, sticking, or penetration. Modified vegetable oils and methylated seed oils are essentially nontoxic to most organisms, including plants. Little is known of the environmental fate of these adjuvants.

Polydimethylsiloxanes are insoluble in water and typically sorb to particulates. Degradation time varies depending on moisture in soils. These chemicals appear to be relatively nontoxic to most organisms, but information is scarce regarding the toxicity and environmental fate of polydimethylsiloxanes. Similarly, little is known about the toxicity or environmental fate of lecithins, which are a commonly used amphoteric surfactant derived from soybeans.

Impact HH-6: Impacts to human health from the use of pesticide adjuvants would be **less than significant** and mitigation is not required.

7.2.6 Biological Control Alternative

Biological control of mosquitoes and other vectors involves the intentional use of vector pathogens (diseases), parasites, and/or predators to reduce the population size of target vectors. Biological control is used as a method of protecting the public from mosquitoes and the diseases they transmit using mosquito parasites, pathogens, and predators. At present, mosquito parasites are not commercially available for mosquito control. The Biological Control Alternative as the District practices it at present would be a continuation of existing activities focused on mosquitofish using applicable techniques, equipment, vehicles, and water craft.

7.2.6.1 Mosquito Larvae Pathogens

As part of their Biological Control Alternative, the District employs bacterial larvicides that are highly specific to mosquitoes. These controls include Bs, Bti, and spinosad. Bs is the only live pathogen being used; Bti and spinosad pesticides are the by-products of bacterial organisms.

All three bacteria are naturally occurring soil organisms, which are commercially produced as mosquito larvicides. Because these forms of biological control are regulated by USEPA and are applied in a similar manner to chemical pesticides, they are evaluated under Section 7.2.7, Chemical Control Alternative, including the discussion of potential impacts. The environmental fate and toxicity of these control agents are described in detail in Appendix B.

7.2.6.2 Mosquito Predators

Mosquitofish (*Gambusia affinis*) are presently the only commercially available mosquito predators. The District's use of these fish in mosquito habitats is the most commonly used biological control agent for mosquitoes in the world. Used correctly, this fish can provide safe, effective, and persistent suppression in various mosquito producing sources. However, due to concerns that mosquitofish may potentially impact red-legged frog and tiger salamander populations, the District limits the public's use of mosquitofish to ornamental fish ponds, water troughs, water gardens, fountains, and unmaintained swimming pools.

Impact HH-7: No impact would occur to human health from the use of mosquitofish. **7.2.6.3 Other Vectors**

No effective natural predators exist to control high rodent populations. Domestic and feral cats may provide short-term control when the rodent population is low, but they can also impact bird populations. The District does not employ cats for rat control. Currently, no commercial biological control agents or products are available for wasp, yellow jacket, and tick control.

7.2.7 Chemical Control Alternative

Chemical control involves the application of nonpersistent selective insecticides to directly reduce populations of mosquitoes and other invertebrate threats to public health and the use of rodenticides to control rats and mice. If and when inspections reveal that mosquitoes or other vector populations are present at levels that trigger the District's criteria for chemical control – based on the vector's abundance/density, species composition, proximity to human settlements, water temperature, or presence of predators – the District applies pesticides to the site in strict accordance with label instructions and federal and state guidelines.

Most of the chemical controls the Program uses are for mosquito control and abatement and are classified as larvicides or adulticides. Below is a discussion of the larvicides, adulticides, and rodenticides the District uses. The active ingredients that were identified as warranting further evaluation in Appendix

B due to their potential toxicity and/or prevalent use/public concern are listed in Table 7-6 (including herbicides discussed previously in Section 7.2.5).

Table 7-6 Active Ingredients Identified for Further Evaluation in Appendix B

Active Ingredient	Vector	Potential Issue
Methoprene	Mosquito	Prevalent use; toxicity to aquatics and insects
Etofenprox	Mosquito	Toxicity to aquatic organisms; no synergist required
Bti	Mosquito	Prevalent use; public concerns
Pyrethrins	Mosquito	Prevalent use; requires synergist (PBO)
Resmethrin	Mosquito	Requires synergist (e.g., PBO); potential endocrine disruptor
Vegetable Oil	Mosquito	Contains low percentage of petroleum distillate
Bromadiolone	Rat	Toxicity to nontarget organisms including mammals, birds, aquatics
Glyphosate	Weeds	Prevalent use; possible endocrine disruptor

7.2.7.1 Mosquito Larvicides

Larvicides are used to manage immature life stages of mosquitoes including larvae and pupae in aquatic habitats. Temporary aquatic habitats are usually targeted because permanent waterbodies generally support natural mosquito predators such as fish. The larvicides are applied using ground application equipment, fixed-wing aircraft, and rotary-wing aircraft. The mosquito larvicides the District uses and where they are discussed in detail in Appendix B are listed in Table 7-7.

Table 7-7 Chemicals Employed for Larval Mosquito Abatement

Chemical Classification	Active Ingredient	Appendix B
Organophosphate	Temephos	Section 4.2.2
Bacterial larvicide	Bs	Section 4.3.1
Bacterial larvicide	Bti	Section 4.3.2
Bacterial larvicide	Spinosad	Section 4.3.3
Hydrocarbon ester	Methoprene	Section 4.3.4
Surfactant	Alcohol Ethoxylated Surfactant (monomolecular film)	Section 4.3.5
Surfactant	Aliphatic Solvent (mineral oil)	Section 4.3.6

7.2.7.1.1 Bacterial Larvicides (Bs, Bti, and Spinosad)

Bacterial larvicides such as Bs and Bti are highly selective microbial pesticides for mosquitoes whose protein spores, when ingested, cause destruction of the gut wall leading to paralysis and death. Another bacterium, *Saacharopolyspora spinosa*, produces spinosyns, which are highly effective mosquito neurotoxicants. All three bacteria are naturally occurring soil organisms and are commercially produced as mosquito larvicides. Unlike Bti and *S. spinosa*, Bs is a live bacterium that can reproduce in natural settings for some time following release. Bs and Bti are applied on a variety of standing and moving waterbodies. The spores of Bs and Bti can persist in the environment for months, but the endotoxins are readily degraded by UV light and persist only for a few hours to a maximum of a few days. Bacterial spores of Bti are uniquely toxic to nematoceran Diptera (mosquitoes, some midges, blackflies, psychodids, and ceratopogonids) (Lacey and Mulla 1990) and do not exhibit any human toxicity.

Spinosad alters nicotine acetylcholine receptors in insects, causing constant involuntary nervous system impacts ultimately leading to paralysis and death. The USEPA has classified spinosad as a “reduced risk” compound because it is an alternative to more toxic, OP insecticides (CDPR 2002). It exhibits acute toxicity to target organisms by all exposure routes but has not been shown to elicit acute or chronic toxicity in humans..

Impact HH-8: No impact would occur to human health from the use of bacterial larvicides.

7.2.7.1.2 Hydrocarbon Ester - Methoprene

Methoprene is an insect growth regulator and selective larvicide. It used in a variety of settings including indoors and outdoors at residences, animal husbandry premises, industrial sites, irrigation systems, and standing waterbodies. It is applied either in response to observed high populations of mosquito larvae at a site, or as a sustained-release product that can persist for 4 months or longer if a site has limited accessibility and has regularly produced immature mosquitoes in the past. It is applied by hand, from an ATV, from watercraft, or from aircraft, including low-flying helicopters (particularly for marshes and other highly vegetated areas) but never when winds exceed 10 mph to prevent drift.

Methoprene has very low acute toxicity to humans and mammals by all routes (USEPA 1991a). No potentially significant impact exists to humans by exposure to typical levels of methoprene. To achieve toxicity to humans, exposures hundreds of times higher than what is legally allowed for use in vector control would be required and such exposure would need to be extensive. Methoprene is of public concern due to its potential ecological effects and widespread use (discussed in Chapter 6, Section 6.2.7.1.2).

Impact HH-9: No impact would occur to human health from the use of the mosquito larvicide methoprene.

7.2.7.1.3 Alcohol Ethoxylated Surfactant (Monomolecular Film)

The monomolecular film formulation used as a surfactant in California for mosquito larvae control is Agnique. Monomolecular films spread a thin film on the water surface that makes it difficult for mosquito larvae, pupae, and emerging adults to attach to the water’s surface, causing them to drown (USEPA 2007a). The films also disrupt larval respiration of some other classes of air-breathing aquatic insects. They are used on an assortment of waterbodies including ornamental ponds, pastures, irrigation systems, drainage systems, and drinking water systems (CDPR 2010a). No evidence supports that these surfactants are toxic to humans.

7.2.7.1.4 Aliphatic Solvents (Mineral Oil)

Monomolecular films are alcohol ethoxylated surfactants (such as BVA-2), which are low-toxicity pesticides that spread a thin film on the surface of water that makes it difficult for mosquito larvae, pupae,

and emerging adults to attach to the water's surface, causing them to drown (USEPA 2007a). The films also disrupt larval respiration of some other classes of air-breathing aquatic insects. They are used on an assortment of waterbodies including ornamental ponds, pastures, irrigation systems, drainage systems, and drinking water systems (CDPR 2010a).

Aliphatic solvents such as mineral oil are applied to water surfaces to form a coating on top of water surfaces to drown larvae, pupae, and emerging adult mosquitoes. They are the product of petroleum distillation and, thus, are complex mixtures of long-chain aliphatic compounds. They are applied to a variety of waterbodies, including swamps, marshes, and intermittently flooded areas (CDPR 2010a). Aliphatic solvents are often used when monomolecular films (alcohol ethoxylated surfactants) do not provide sufficient mosquito control. They also break down more rapidly (2 to 3 days) and are practically nontoxic to most nontarget organisms. They have a low degree of acute toxicity to mammals. Therefore, mineral oil should not result in adverse effects to human health when applied using District BMPs.

Impact HH-10: No impact would occur to human health from the use of alcohol ethoxylated and aliphatic solvent surfactant larvicides.

7.2.7.1.5 Temephos

Temephos is the only OP with larvicidal use and is used to help prevent mosquitoes from developing resistance to the bacterial larvicides. It is the only OP included in the District's Program. It was used prevalently in California for mosquito abatement from 1965 into the mid-1980s; however, microbial pesticides (e.g., Bs, Bti, spinosad), methoprene, and surface oils are used much more frequently now. It is used in various waterbodies including lakes, marshes, drainage systems, irrigation systems, and polluted and stagnant water; it is not used on agricultural lands (CDPR 2010a). The District primarily applies temephos to man-made sources such as tire piles, utility vaults, and cemetery urns. Temephos has extremely low water solubility and binds strongly to soils. It has low toxicity for vertebrates at the levels used for mosquito control (USEPA 2000). The USEPA (2000) states that people are likely not exposed to temephos in drinking water or from residential use.

Impact HH-11: Impacts to human health from the use of temephos would be **less than significant** and mitigation is not required.

7.2.7.2 Mosquito Adulticides

In addition to chemical control of mosquito larvae, the District may use pesticides for control of adult mosquitoes when no other tools are available and if specific criteria are met, including species composition, population density (as measured by landing count or other quantitative method), proximity to human populations, and/or human disease risk. Adulticide materials are used infrequently and only when necessary to control mosquito populations (e.g., those areas with treeholes where access to larval breeding sites is impractical).

Adulticides the District potentially uses include pyrethrins, synthetic pyrethroids, pyrethroid-like compounds, OPs, and synergists. Table 7-8 lists the adulticides the District uses for vector abatement. Several of these active ingredients, as well as a few others, are also used for the control of yellow jacket wasps and, in some cases, to control tick populations that pose an imminent threat to people, pets or livestock (Table 7-8 and this section). A subset of these active ingredients required further evaluation in Appendix B and further discussion is provided below. A detailed discussion of the environmental fate and toxicity of these pesticides is provided in Appendix B.

Table 7-8 Chemicals Employed for Adult Insect Abatement

Chemical Classification	Active Ingredient	Vector	Appendix B
Pyrethrin	Pyrethrins	Mosquito; yellow jacket	Section 4.1.1
Pyrethroid	Resmethrin	Mosquito	Section 4.1.2
Pyrethroid	Permethrin	Mosquito	Section 4.1.3
Pyrethroid	Etofenprox	Mosquito	Section 4.1.4
Synergist	PBO	Mosquito; yellow jacket	Section 4.1.5
Organophosphate	Naled	Mosquito	Section 4.2.1

7.2.7.2.1 Pyrethrins

Pyrethrins are naturally occurring compounds the flowers of the *Chrysanthemum* species produce. They effectively induce temporary paralysis in insects but are not acutely lethal by themselves; thus, they are used concomitantly with the synergist PBO, which inhibits metabolism of the pyrethrins so that a lethal dose is assured (USEPA 2006c). The District uses pyrethrins on crops, animal husbandry premises and pastures, outdoor household areas, and for wide-area mosquito abatement in areas that include aquatic habitats.

Pyrethrins have low to moderate acute mammalian toxicity via the oral, dermal, and inhalation routes (Categories III and IV). They are a moderate eye irritant (Category III), a mild dermal irritant (Category IV), and not a skin sensitizer. The effects of pyrethrins are (1) neurobehavioral effects following acute, short-term, and chronic exposure, with nervous system lesions observed in the rat and mouse following acute exposure; (2) thyroid effects, following chronic exposure in the rat and dog; and (3) liver effects, following short- and long-term exposure in the rat, dog, and mouse. The neurobehavioral effects are considered relevant to humans because the effects are observed in both the rat and mouse, and the mode of action affects a basic function of the nervous system that is common to all animals (USEPA 2006c).

They are of concern because they are used prevalently and require the use of the synergist PBO, a potential endocrine disruptor (USEPA 2009a). However, the District uses pyrethrins only when absolutely necessary and in minimal amounts in ULV applications that are designed to break down rapidly, resulting in very low potential exposure to humans.

Impact HH-12: Impacts to human health from the use of pyrethrins would be **less than significant** and mitigation is not required.

7.2.7.2.2 Pyrethroids, Pyrethroid-Like Compounds, and Synergists

Pyrethroids are synthetic compounds that are chemically similar to the pyrethrins but have been modified to increase stability and activity against insects. Pyrethroids bind to neuronal voltage-gated sodium channels, preventing them from closing; this persistent activation of the channels then leads to paralysis. First generation or "Type I" pyrethroids include d-allethrin, phenothrin (sumithrin), resmethrin, and tetramethrin. These pyrethroids are used to control flying and crawling insects in a number of commercial and horticultural applications and are sold for residential use and application on pets to control fleas and ticks. They have effective insect knockdown capabilities but are unstable as they are highly photosensitive (i.e., easily degraded by light). The newer second-generation/"Type II" pyrethroids contain an α -cyano group, which reduces their photosensitivity, thereby increasing their persistence and toxicity. The active ingredients that fall into this group include deltamethrin, esfenvalerate, lambda-cyhalothrin, and permethrin.

Some synthetic insecticides are similar to pyrethroids, such as etofenprox, but have a slightly different chemical composition. The pyrethroids that were identified for further evaluation in Appendix B are discussed below.

7.2.7.2.3 Resmethrin

Resmethrin is the active ingredient in Scourge®. It is a restricted-use pesticide due to its toxicity to fish and is available for use only by certified pesticide applicators or persons under their direct supervision. Resmethrin has low acute toxicity via the oral (Category III), dermal (Category III), and inhalation (Category IV) routes of exposure. Resmethrin is included in the final list of chemicals for screening under USEPA's Endocrine Disruptor Screening Program (USEPA 2009a).

Public concern regarding the potential effects of resmethrin has severely restricted use of this ingredient, primarily because of the possible effects it may have as an endocrine disruptor., Scourge® is rarely used, if at all, and is being phased out of the District's Program and replaced with a nonresmethrin alternative.

7.2.7.2.4 Permethrin

Permethrin is also a pyrethroid. Dermal exposure in humans can cause tingling and pruritus with blotchy erythema on exposed skin (ATSDR 2003). In humans, acute effects observed subsequent to ingestion of permethrin included nausea, vomiting, abdominal pain, headache, dizziness, anorexia, and hypersalivation. Reports of severe poisoning are rare and usually follow ingestion of substantial, but poorly described, amounts of permethrin. Symptoms of severe poisoning include impaired consciousness, muscle fasciculation, convulsions, and noncardiogenic pulmonary edema (ATSDR 2003). Systemic effects are similar to those seen in acute and chronic ingestion with prolonged contact or contact with high concentrations of permethrin. Acute toxicity to permethrin via inhalation has been shown to be very small. The USEPA (2006c) has classified permethrin as Category III for acute oral and acute dermal toxicity, Category III for eye irritation potential, and Category IV for dermal irritation potential. Although often used for residential insect control, products containing permethrin are considered very low risk to humans. Acute, chronic noncancer, and cancer dietary risks from permethrin have been shown to be below the USEPA's level of concern. Occupational (handler/applicator) uses are also below USEPA noncancer levels of concern for typical uses, and typical personal protective equipment may be used.

Because permethrin is included in the final list of chemicals for screening under USEPA's Endocrine Disruptor Screening Program (USEPA 2009a), it is of concern to the public. However, the District rarely uses it, applies it through ULV application, or using a backpack mister or a handcan/duster, and does not apply it during high winds.

7.2.7.2.5 Etofenprox

Etofenprox is a pyrethroid-like insecticide that is the active ingredient in Zenivex. It differs in structure from pyrethroids in that it lacks a carbonyl group and has an ether moiety, whereas pyrethroids contain ester moieties. It is used indoors, as a spot treatment for pets, and as an outdoor fogger to control flying and crawling insect pests. It is frequently applied to backyards and patios and sometimes directly to domestic pets. As with many other pyrethroids, etofenprox has low acute toxicity to humans and other mammals and has been shown to be below the USEPA level of concern under appropriate uses. The public's concerns regarding the ecological impacts of etofenprox are primarily due to the concern that it may be an endocrine disruptor. It is further discussed in Chapter 6, Section 6.2.7.2.5.

Impact HH-13: Impacts to human health from the use of pyrethroids and pyrethroid-like compounds as mosquito adulticides would be **less than significant** and mitigation is not required.

7.2.7.2.6 Piperonyl Butoxide

PBO is a pesticide synergist that enhances the effectiveness of pesticide active ingredients, such as pyrethrins and pyrethroids, by inhibiting microsomal enzymes and, thus, the breakdown of the other active

ingredient(s) (USEPA 2006b). It is a registered active ingredient in products used to control flying and crawling insects and arthropods in agricultural, residential, commercial, industrial, and public health settings. No products contain only PBO. It degrades quickly in soil and water. PBO has a low acute toxicity by oral, inhalation, and dermal routes, but it is included in the final list of chemicals for screening under USEPA's Endocrine Disruptor Screening Program (USEPA 2009a). As a synergist, PBO is applied using the same guidelines as those for pyrethroids and pyrethrins: ULV application, a backpack mister, or handcan/duster, and it is not applied during high winds.

Impact HH-14: Impacts to human health from the use of the synergist PBO in mosquito adulticides would be **less than significant** and mitigation is not required.

7.2.7.3 Yellow Jacket Wasp and Tick Adulticides

The District selectively applies insecticides to control ground-nesting yellow jackets that pose an imminent threat to people or pets. This activity is generally triggered by public requests for District assistance or action rather than as a result of regular surveillance of their populations. The District does not apply pesticides to yellow jacket populations that are located in or on a structure. Whenever the District learns that a hive is situated in or on a structure or is above ground, the resident(s) are encouraged to contact a private pest control company that is licensed to treat the infestation. Yellow jacket nests that are off the ground would be treated only under special circumstances to protect public health and safety of the District's residents. When a District technician encounters a honeybee swarm or unwanted hive, residents are referred to the County Agricultural Commissioner's Office, which maintains a referral list of beekeepers that can safely remove the bees. If a District technician deems it appropriate to treat stinging insects, they will apply the insecticide directly within the nest in accordance with the District's policies to avoid drift of the insecticide or harm to other organisms. Alternatively, they will place tamper-resistant traps or bait stations, selective for the target insect in the immediate environment of the vector.

Tick populations can be managed somewhat with pyrethroids but widespread area treatments are not practical because host animals tend to reintroduce them. Tick management for the District is limited at present to collection, identification, and testing for certain tickborne diseases; and information is provided to the public about their biology and natural ways to minimize interactions with this vector organism.

Pyrethroid-based chemicals are typically used against ground-nesting yellow jackets and are applied directly into the underground nest, which prevents drift and further reduces the potential for nontarget exposure to these compounds. In addition to the pyrethrins and pyrethroids discussed above, the District uses lambda-cyhalothrin to control yellow jacket wasps.

7.2.7.4 Rodenticides

The District employs limited use of rodenticides in response to District resident requests. The District may use two different groups of anticoagulant rodenticides, known as first generation and second generation rodenticides. First generation rodenticides require consecutive multiple doses or feedings over a number of days to be effective. Second-generation rodenticides require only one dose and are effective against rodents that have become resistant to first generation rodenticides; they are far more toxic than first generation baits.

The District may conduct rodent baiting at underground sites such as sewers, storm drains, or catch basins. Secure bait stations or other accepted methods of rodent baiting are conducted in areas with severe rodent infestations. In sewer baiting, bait blocks containing a rodenticide are suspended by wire above the water line to encourage rodent feeding. The rodenticides the District uses, proposes to use, and where they may be used are listed in Table 7-9 and are discussed in detail in Appendix B in the section indicated.

Table 7-9 Chemicals Employed for Rodent Abatement

Chemical Classification	Active Ingredient	Appendix B
First-generation anticoagulant	Diphacinone	4.5.1
Second-generation anticoagulant	Bromadiolone	4.5.2
Second-generation anticoagulant	Difethialone	4.5.3
Sterol	Cholecalciferol	4.5.4

7.2.7.4.1 Anticoagulant Rodenticides

As their name suggests, anticoagulants function by inhibiting the production of blood-clotting factors. First-generation (e.g., diphacinone) anticoagulants require rodents to consume the bait for several consecutive feedings for delivery of a lethal dose. Diphacinone baits are used around buildings and similar man-made structures. These compounds have very low water solubility and are moderately persistent in soils. First-generation rodenticides are classified as Category I (highly toxic) to mammals for oral, dermal, and inhalation toxicity (USEPA 1998b). As bait poisons, these anticoagulants are used sparingly as the chance of nontarget animals consuming the bait is a possible unwanted nontarget effect. For this reason, the baits are made to be less accessible to other animals by special packaging.

Second-generation (e.g., brodifacoum, bromadiolone) compounds have the same mode of action as first-generation anticoagulants but are more acutely toxic than the first-generation anticoagulants. The second-generation anticoagulant rodenticides are substantially more potent than the first-generation compounds, and a lethal dose can be ingested in a single feeding. Because they are retained much longer in body tissues, second generation compounds are much more likely to poison predatory wildlife or pets that eat live or dead poisoned prey (Hartless and Jones 2011). Brodifacoum has the greatest acute toxicity of the Program rodenticides, but it is used very infrequently.

Products containing second-generation active ingredients are no longer permitted to be sold to the general public. These products remain available to professional pest control personnel, and strict adherence to product label requirements and District BMPs can ensure their safe use for controlling and eradicating nuisance rodent populations, including the use of tamper-proof bait stations; securing bait stations at deployment locations to prevent disruption and/or removal by wildlife; and proper education of citizens including residents about the potential risk to pets, wildlife, and children.

Impact HH-15: Impacts to human health from the use of anticoagulant rodenticides would be **less than significant** and mitigation is not required.

The anticoagulant rodenticide bromadiolone was selected for further evaluation in Appendix B and is discussed individually below.

7.2.7.4.2 Bromadiolone

The District does not currently use bromadiolone, the active ingredient in Contrac products, but it is a commercially available second-generation anticoagulant rodenticide that is marketed for use around buildings and in transport containers, alleys, and sewers. It is highly toxic to mammals, including humans, by acute oral, dermal, and inhalation exposure (USEPA 1998b). Bromadiolone is listed as Category III for eye irritation and Category IV for skin dermal irritation (USEPA 1998b).

Bromadiolone is a concern to the public due to its high mammalian toxicity. However, bromadiolone is usually wax-encased (e.g., Contrac Blox) in block form, which has exceptionally low water solubility and low leaching potential. Furthermore, when the District applies bromadiolone blocks in sewers, usually below manhole covers, it is suspended by a wire. This method of bait deployment reduces the probability of exposure to humans and pets. When bromadiolone is used around residences, usually at a

homeowner's request, the District places bromadiolone in tamper-proof bait stations, which are anchored at treatment locations (e.g., wires, stakes) to ensure that they cannot be dragged away by children or pets. The District educates citizens about the locations of bait blocks and potential risks to pets and children.

Impact HH-16: Impacts to human health from the use of bromadiolone would be **less than significant** and mitigation is not required.

7.2.7.4.3 Difethialone

Difethialone is also a second-generation anticoagulant rodenticide that is the active ingredient in FirstStrike Soft Bait. It is very toxic to mammals, including humans, domestic pets, and nontarget mammalian wildlife, by all acute exposure routes. Difethialone is not known to cause skin or eye irritation. No genotoxic or carcinogenic effects have been noted (Annex I - Norway 2007).

Difethialone is also of public concern because of its high mammalian toxicity and because it is used in areas frequented by humans and domestic animals (parks, landscaped areas) during much of the year. As with bromadiolone, the District uses difethialone in tamper-proof bait stations, which are anchored at treatment locations (e.g., wires, stakes) to ensure that they cannot be dragged away by children or pets. The rodenticides are used at historical baiting sites when food competition occurs to increase the likelihood of exposure to only target rodents. The District educates citizens about the locations of bait blocks and potential risks to pets and children. If the District expands use in the future or additional issues arise regarding the use of this rodenticide, new, more protective rodenticide bait station alternatives the USEPA reports could be considered (<http://www.epa.gov/pesticides/mice-and-rats/rodent-bait-station.html>).

Impact HH-17: Impacts to human health from the use of difethialone would be **less than significant** and mitigation is not required.

7.2.7.4.4 Sterol – Cholecalciferol

Cholecalciferol is used in and around homes, industrial buildings, and similar man-made structures, agricultural buildings, transport vehicles, port and terminal buildings; and in alleys. Formulation types include pellets and blocks (Clock-Rust and Sutton 2011). It is essentially insoluble in water and immobile in soils. Cholecalciferol is vitamin D3, and ingestion of high amounts results in hypercalcemia from mobilization of calcium from bone matrix into blood plasma leading to soft tissue mineralization resulting in loss of functionality of kidneys, cardiac muscle, etc. (Morrow 2001). It is not acutely toxic to humans at the doses used in bait stations and the District uses it infrequently. When the District does use it, residential treatments involve bait station deployment within 50 feet of home.

Impact HH-18: Impacts to human health from the use of cholecalciferol would be **less than significant** and mitigation is not required.

7.2.8 Other Nonchemical Control/Trapping Alternative

The trapping of rodents and/or yellow jackets is conducted on a limited basis when these organisms pose a threat to public health and welfare. For both vector species, District staff place the tamper-resistant or baited trap(s) primarily at the request of the property owner or manager. The District does not remove rats or yellow jackets that are in or on structures. When these structural requests are made, residents are referred to a directory of private pest control companies. The use of traps is not an effective method for managing yellow jacket populations, and in some circumstances may not be very effective when dealing with large populations of rodents. Trapping is useful when collecting rodents for disease testing or to better estimate population levels of yellow jackets and rodents but is not a practical vector control option for control of these vector populations. No impact to human health is expected from the District's use of this alternative.

Impact HH-19: No impact would occur to human health from the District's use of the Other Nonchemical Control/Trapping Alternative.

7.2.9 Cumulative Impacts

"Cumulative impacts" are defined as "two or more individual effects which, when considered together, are considerable or compound or increase other environmental impacts (CEQA Guidelines, Section 15355). Cumulative impacts, as they relate to human health, include past, present, and reasonably foreseeable actions that potentially impact humans. Cumulative impacts can result from individually minor, but collectively significant, projects taking place over a period of time. The cumulative impact analysis is contained in Section 13.5 and focuses on the potential for the use of pesticides for mosquito and vector control to contribute to regional pesticide use, which is of concern for its potential impacts to the health of human populations. It includes Table 13-1, Historical Pesticide Use within the NCMAD Program Area and Table 13-2, Pesticide Use by NCMAD within the District Service Area.

Although large uncertainty and high variation exist in the reported amounts of pesticide use within the District's Program Area counties, they vary according to particular needs, majority of habitat type, and seasonal vector outbreaks. The public is aware of these pesticide uses and, in general, is pressuring agencies within these counties to use less pesticide whenever possible. The District uses BMPs in their pesticide applications for mosquito and vector control and is attempting to reduce total pesticide use where possible and consistent with IPM practices.

The District's small incremental contributions to overall pesticide use within its Program Area do not trigger a cumulatively considerable impact on pesticide use. While overall use of pesticides throughout the Program Area may be considered cumulatively significant, the District's small incremental contributions to this impact are not cumulatively significant. Therefore, the **Program's long-term activities including chemical applications would not contribute considerably to human health impacts**. The Program alternatives would not result in significant cumulative impacts to the human health condition of the region.

7.2.10 Environmental Impacts Summary

Table 7-10 presents a summary of human health impacts associated with the six alternatives. The human health impacts correspond to those in Sections 7.2.3 through 7.2.8. All of the impacts were determined to be either "no impact" or a "less-than-significant impact."

This page left intentionally blank

Table 7-10 Summary of Alternative Human Health Impacts

Impact Statement	Surveillance	Physical Control	Vegetation Management	Biological Control	Chemical Control	Other Nonchemical/ Trapping
Effects on Human Health						
Impact HH-1: No impact would occur to human health from the use of the Surveillance Alternative.	N	na	na	na	na	na
Impact HH-2: Impacts to human health from use of the Physical Control Alternative would be less than significant and mitigation is not required.	na	LS	na	na	na	na
Impact HH-3: No impact would occur to human health from the nonherbicide Vegetation Management Alternative	na	na	N	na	na	na
Impact HH-4: Impacts to human health from the use of 2,4 D would be less than significant and mitigation is not required.	na	na	LS	na	na	na
Impact HH-5: Impacts to human health from the use of glyphosate would be less than significant and mitigation is not required.	na	na	LS	na	na	na
Impact HH-6: Impacts to human health from the use of pesticide adjuvants would be less than significant and mitigation is not required.	na	na	LS	na	na	na
Impact HH-7: No impact would occur to human health from the use of mosquitofish.	na	na	na	N	na	na
Impact HH-8: No impact would occur to human health from the use of bacterial larvicides.	na	na	na	na	N	na
Impact HH-9: No impact would occur to human health from the use of the mosquito larvicide methoprene.	na	na	na	na	N	na
Impact HH-10: No impact would occur to human health from the use of alcohol ethoxylated and aliphatic solvent surfactant larvicides.	na	na	na	na	N	na
Impact HH-11: Impacts to human health from the use of temephos would be less than significant and mitigation is not required.	na	na	na	na	LS	na
Impact HH-12: Impacts to human health from the use of pyrethrins would be less than significant and mitigation is not required.	na	na	na	na	LS	na
Impact HH-13: Impacts to human health from the use of pyrethroids and pyrethroid-like compounds as mosquito adulticides would be less than significant and mitigation is not required.	na	na	na	na	LS	na

Table 7-10 Summary of Alternative Human Health Impacts

Impact Statement	Surveillance	Physical Control	Vegetation Management	Biological Control	Chemical Control	Other Nonchemical/ Trapping
Impact HH-14: Impacts to human health from the use of the synergist PBO in mosquito adulticides would be less than significant and mitigation is not required.	na	na	na	na	LS	na
Impact HH-15: Impacts to human health from the use of anticoagulant rodenticides would be less than significant and mitigation is not required.	na	na	na	na	LS	na
Impact HH-16: Impacts to human health from the use of bromadiolone would be less than significant and mitigation is not required.	na	na	na	na	LS	na
Impact HH-17: Impacts to human health from the use of difethialone would be less than significant and mitigation is not required.	na	na	na	na	LS	na
Impact HH-18: Impacts to human health from the use of cholecalciferol would be less than significant and mitigation is not required.	na	na	na	na	LS	na
Impact HH-19: No impact would occur to human health from the District's use of the Other Nonchemical Control/Trapping Alternative.	na	na	na	na	na	N

LS = Less-than-significant impact

N = No impact

na = Not applicable

SM = Potentially significant but mitigable impact

SU = Significant and unavoidable impact

7.2.11 Mitigation and Monitoring

All impacts to human health are identified as either “no impact” or a “less-than-significant impact.” Therefore, mitigation measures are not required due to the insignificant impacts identified for all of the Program alternatives described.

This page left intentionally blank